Alpine ecosystems under global change

Auswirkungen langfristiger Umweltveränderungen auf alpine Ökosysteme

Christian Körner und Erika Hiltbrunner

Department of Environmental Sciences
University of Basel, Switzerland

Tagung Parkforschung Schweiz, 29.Oktober 2019

Nival

Snowline

Alpine

Uplands

Treeline

Montane

Lowlands

© Ch Körner

Alexander von Humboldt‘s 250th birthday

→ Humboldt invented the isotherm concept



Elevation *per se* does not matter. What matters is climate

Humboldt's ‚Naturama' redesigned: the wold is connected through mountains

The alpine and montane belts globally



Körner C (2003) Alpine plant life. Springer, Berlin

Mountains are more biodiverse than expected by their area

Number of named species at 2500 m,
350 m above treeline, across 1 km^2

554 Insects (incl. chilopods)
313 Fungi (Basidio-, Asco-, Glomeromycota)
304 Flowering plants
300 Lichens
215 Diatoms
166 Mosses
128 Spiders and Mites
30 Vertebrates (birds, rodents, amphibia, reptiles)

In total 2098 species

excluding
Bacteria
Nematodes
Collembols

Hotspot Furka
Hiltbrunner E, Körner C (2018)

Climatic warming opens new land


fundamental niche
realized niche

No question, alpine organisms do in large not experience the climate weather stations report, but trees do ...



That is why the treeline is a perfect reference for defining life zones in mountains







C Körner (2007) Erdkunde 61:316

©Ch Körner

Habitat diversity is the regional driver of biodiversity

→ a safeguard against species loss



© Ch Körner

Microtopogaphic
diversity facilitates
species diversity

Habitat loss due to climate warming

(assuming proportionality between air temperature and plant temperature)



Scherrer D & Körner C (2011) J Biogeogr 38:406

© Ch Körner

Species responses to climatic warming

Mountains may be refugia (2, 4)

traps (3, 5)

or a chance (6)



... but some habitats will shrink in size

©Ch Körner

Projections of longterm effects of climate warming depend on spatial resolution

Surface temperature (IR) bright days and daylight hours only

Spatial resolution (m^2)	2 K scenario habitat loss (%)	3 K scenario habitat loss (%)	4 K scenario habitat loss (%)
1 x 1	7.5 ± 0.9	20.3 ± 8.0	40.4 ± 19.0
5 x 5	16.8 ± 9.9	40.0 ± 22.2	57.5 ± 30.0
10 x 10	17.4 ± 7.6	42.8 ± 20.1	67.1 ± 24.3
25 x 25	44.1 ± 24.3	68.1 ± 25.6	86.2 ± 11.3
100 x 100	64.7 ± 28.9	95.5 ± 3.7	100 ± 0.0

Means for three different slopes at 2200 - 3000 m of elevation in the Alps

Scherrer D et al. (2011) J Biometeorology, see also CF Randin *et al* (2009) GCB 15:1557

© Ch Körner


dangerous
safe
Phenology
© Ch Körner


Phenology ensures sexual reproduction and survival
Absolute frequency (n=42 species)
0
10
20
30
40
vegetative
bud visible
first
full
end flower
mature seeds/fruits
first
seeds/fruits released
10.6.
17.6.
26.6.
30.6.
1.7.
6.7.
7.7.
14.7.
16.7.
19.7.
20.7.
23.7.
25.7.
29.7.
31.7.
5.8.
7.8.
20.8.
26.8.
1.10.
Flowering phenology at 2440 m in 2015

Half of the high alpine/nival taxa are 'opportunistic‘, the other half is strongly controlled by photoperiodism

There are options for selection: later flowering

Damage between -3 and -12 ° C (summer)
depending on species and tissue type

Heat waves can induce drought even in alpine plants


Sharp snow-melt gradients:
range limits at small scale
23 July, 2016
16 July, 2016




Alpine LTER sites - Long Term Ecological Monitoring
48°
46°
44°
Nationalpark
Hohe Tauern
Südtirol-EURAC
Furka-ALPFOR
0
100
km
4000 m
3500 m
3000 m
2500 m
2000 m
1500 m
1000 m
500 m
0 m

Top of transect

Bottom of transect

Biomass across snow melt gradient in 2018



Peak season plant biomass at 2300 – 2500 m asl
Comparison between 2017 and 2018



Plant biomass production along a snow-melt gradient in summer 2017, Furka-Pass, Switzerland, 2467 m a.s.l.





Vertical Biomass allocation along a snow melt gradient



The problem of year to year variation of productivity in long-term observations with too large census intervals.
Conclusions B and C differ strongly from the trend in A.

Elevated CO_2: No growth stimulation in 4 years in alpine grassland (2500 m)



Körner C *et al* (1997) Acta Oecol 18:165

Glacier forefield plants:
No growth stimulation by eCO_2 in 3 years despite NPK-addition, some species grew even less
N Inauen *et al* (2012) GCB 18:985
©Ch Körner

The effect of N deposition on alpine plants

Swiss Alps, 2500 - 2600 m, critical load 5 -10 kg N ha^{-1} a^{-1}



C Heer & Ch Körner (2002) Basic Appl Ecol 3:39

E Hiltbrunner, unpubl data

©Ch Körner

Furka
N-Deposition, both from NW and SE (the Po-Region)
NASA

Nitrogen deposition: winners and losers

E. Hiltbrunner, Basel


Biomass effects at >5 kg N ha-1 a-1 after 4 years
(E. Hiltbrunner, Basel, unpubl.)
Biomasse 2007
(kg m-2)
0.4
0.3
0.2
0.1
0.0
**
**
0
2.5
5
10
15
20
25
kg N ha-1 a-1
©Ch Körner

Summary

- Alpine plants, ‘engineer‘ their microclimate
- 10K contrast of seasonal mean T over few m
- Phenology x snow cover x freezing matters
- Alpine plants are not carbon limited
- N deposition is more influential than changes in mean temperature
- Most alpine plants not more ‘vulnerable‘

Alpine Plant Ecology

Summer School on Alpine Plant Life

Swiss central Alps, 12- 18 July 2020

Erika Hiltbrunner, Christian Körner, Jürg Stöcklin
University of Basel
www.alpfor.ch

Location:
ALPFOR
Alpine Research and Education Station
Furka, 2440 m a.s.l., Swiss central Alps

Registration: PhD students register at franziska.grob@unibas.ch, at MOnA (Univ. of Basel students) and PSC PhDs at PSC. Pre-registration (with motivation letter) until 28 Feb. 2020. Acceptance information: 3 April 2020, confirmed registration: 29 May 2020.

‘Experiments‘ by nature: opportunities for basic research and conservation