



Nitrogen but not phosphorus addition affects symbiotic N₂ fixation by legumes in natural and semi-natural grasslands located on four continents

Eduardo Vázquez¹ · Per-Marten Schleuss² · Elizabeth T. Borer³ · Miguel N. Bugalho⁴ · Maria C. Caldeira⁵ · Nico Eisenhauer⁶ · Anu Eskelinen⁷ · Philip A. Fay⁸ · Sylvia Haider⁹ · Anke Jentsch¹⁰ · Kevin P. Kirkman¹¹ · Rebecca L. McCulley¹² · Pablo L. Peri¹³ · Jodi Price¹⁴ · Anna E. Richards¹⁵ · Anita C. Risch¹⁶ · Christiane Roscher¹⁷ · Martin Schütz¹⁸ · Eric W. Seabloom¹⁹ · Rachel J. Standish²⁰ · Carly J. Stevens²¹ · Michelle J. Tedder²² · Risto Virtanen²³ · Marie Spohn²⁴

Received: 26 January 2022 / Accepted: 16 May 2022
© The Author(s) 2022

Abstract

Background and aims The amount of nitrogen (N) derived from symbiotic N₂ fixation by legumes in grasslands might be affected by anthropogenic N and phosphorus (P) inputs, but the underlying mechanisms are not known.

Methods We evaluated symbiotic N₂ fixation in 17 natural and semi-natural grasslands on four continents

that are subjected to the same full-factorial N and P addition experiment, using the ¹⁵N natural abundance method.

Results N as well as combined N and P (NP) addition reduced aboveground legume biomass by 65% and 45%, respectively, compared to the control, whereas P addition had no significant impact. Addition of N and/or P had no significant effect on the symbiotic N₂ fixation per unit legume biomass. In consequence, the amount of N fixed annually per grassland area was less than half in the N addition treatments compared to control and P addition, irrespective of whether the dominant legumes were annuals or perennials.

Responsible Editor: Euan K. James.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s11104-022-05498-y>.

E. Vázquez (✉) · P.-M. Schleuss · M. Spohn
Department of Soil Ecology, Bayreuth Center of Ecology and Environmental Research (BayCEER), University of Bayreuth, Dr.-Hans-Frisch-Straße 1-3, 95448 Bayreuth, Germany
e-mail: eduardo.vazquez.garcia@slu.se

E. Vázquez · M. Spohn
Department of Soil and Environment, Swedish University of Agricultural Sciences (SLU), Lennart Hjelm's väg 9, 75007 Uppsala, Sweden

E. T. Borer · E. W. Seabloom
Dept. of Ecology, Evolution, and Behavior, University of Minnesota, St. Paul, MN 55108, USA

M. N. Bugalho
Centre for Applied Ecology “Prof. Baeta Neves” (CEABN-InBIO), School of Agriculture, University of Lisbon, 1349-017 Lisbon, Portugal

M. C. Caldeira
Forest Research Centre, School of Agriculture, University of Lisbon, 1349-017 Lisbon, Portugal

N. Eisenhauer · A. Eskelinen · S. Haider · C. Roscher
German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Puschstrasse 4, 04103 Leipzig, Germany

N. Eisenhauer
Institute of Biology, Leipzig University, Puschstrasse 4, 04103 Leipzig, Germany

A. Eskelinen · C. Roscher
Physiological Diversity, Helmholtz Centre for Environmental Research (UFZ), Permoserstrasse 15, 04318 Leipzig, Germany

Conclusion Our results reveal that N addition mainly impacts symbiotic N₂ fixation via reduced biomass of legumes rather than changes in N₂ fixation per unit legume biomass. The results show that soil N enrichment by anthropogenic activities significantly reduces N₂ fixation in grasslands, and these effects cannot be reversed by additional P amendment.

Keywords Grasslands · Legumes · Nitrogen addition · Nutrient Network (NutNet) · Phosphorus addition · ¹⁵N natural abundance method

Introduction

Grasslands cover approximately 40% of the terrestrial ice-free surface of the Earth and provide diverse ecosystem services including climate regulation, soil carbon storage, plant diversity maintenance and support for pollinators while contributing to human nutrition (Lamarque et al. 2011). In particular, legumes are one of the key plant functional groups in grasslands for their capacity to increase the nitrogen (N) availability by symbiotic N₂ fixation, which, in turn, enhances the grassland net primary productivity, mitigates environmental pollution, and increases forage quality

and productivity, critical for livestock production and stockless organic cropping systems (Lüscher et al. 2014; Suter et al. 2015). However, anthropogenic N inputs (in the form of fertilizers, manure, and atmospheric deposition) are changing the supply of N relative to phosphorus (P) in grasslands (Peñuelas et al. 2013) which can affect the symbiotic N₂ fixation by legumes (Høgh-Jensen et al. 2002; Carlsson and Huss-Danell 2003; Stevens et al. 2004). The N and P availability can affect symbiotic N₂ fixation by changing the legume biomass production in grasslands and the contribution of N derived from symbiotic N₂ fixation to the total N content of legumes (percentage of legume N derived from atmosphere, %Ndfa) (Høgh-Jensen et al. 2002; Nyfeler et al. 2011; Peoples et al. 2012). Therefore, understanding the effects of N and P inputs on symbiotic N₂ fixation by legumes is crucial to maintain grassland herbage productivity and additional ecosystem services such as plant biodiversity.

Nitrogen inputs can affect the legume biomass production and the %Ndfa and thus, symbiotic N₂ fixation by legumes in grasslands (West et al. 2005; Nyfeler et al. 2011; Oberson et al. 2013). The %Ndfa often declines with availability of both ammonium and nitrate in soil (Leidi and Rodríguez-Navarro

A. Eskelinen · R. Virtanen
Ecology & Genetics, University of Oulu, PO Box 3000,
90014 Oulu, Finland

P. A. Fay
USDA-ARS Grassland Soil and Water Research
Laboratory, Temple, TX 76502, USA

S. Haider
Institute of Biology / Geobotany and Botanical Garden,
Martin Luther University Halle-Wittenberg, Am Kirchtor
1, 06108 Halle, Germany

A. Jentsch
Disturbance Ecology, Bayreuth Center of Ecology
and Environmental Research (BayCEER), University
of Bayreuth, Universitaetsstr. 30, 95447 Bayreuth,
Germany

K. P. Kirkman · M. J. Tedder
School of Life Sciences, University of KwaZulu-Natal,
Carbis Road, Pietermaritzburg 3209, South Africa

R. L. McCulley
Department of Plant and Soil Sciences, University
of Kentucky, Lexington, KY 40546, USA

P. L. Peri
Instituto Nacional de Tecnología Agropecuaria (INTA),
Universidad Nacional de La Patagonia Austral (UNPA) -
CONICET CC 332, CP 9400 Río Gallegos, Santa Cruz,
Argentina

J. Price
Institute for Land, Water and Society, Charles Sturt
University, Albury, NSW 2640, Australia

A. E. Richards
CSIRO Land and Water, 564 Vanderlin Drive, Berrimah,
Northern Territory 0828, Australia

A. C. Risch · M. Schütz
Swiss Federal Institute for Forest, Snow and Landscape
Research WSL, Zuercherstrasse 111, 8903 Birmensdorf,
Switzerland

R. J. Standish
Harry Butler Institute, Murdoch University, 90 South
Street, Murdoch, WA 6150, Australia

C. J. Stevens
Lancaster Environment Centre, Lancaster University,
Lancaster LA1 4YQ, UK

2000; Peoples et al. 2012), because symbiotic N_2 fixation is energetically expensive, and legumes take up reactive N, if available in the soil. In addition, grasses and non-leguminous forbs often displace legumes at high N availability due to their higher competitiveness for light (Soussana and Tallec 2010; Tognetti et al. 2021). Thus, anthropogenic N inputs to grasslands, can lead to decreased N_2 fixation by legumes affecting the grassland functioning.

Symbiotic N_2 fixation by legumes in grasslands can also be influenced by P inputs (Høgh-Jensen et al. 2002; Edwards et al. 2006). Low soil P levels reduce the activity of N_2 -fixing legume-associated bacteria (Edwards et al. 2006) due to the high ATP requirements of N_2 fixation (Valentine et al. 2017). At the plant level, long-term P deprivation decreases nodular P concentration and reduces the energy status of the nodules and their capacity to fix N_2 , shifting the source of legume N nutrition from atmospheric N_2 fixation towards uptake of reactive soil N (Valentine et al. 2017). Low P availability could reduce the rate of N_2 fixation per unit legume biomass, removing the advantage that legumes might have over non- N_2 fixing plants in N-poor conditions and reducing legume biomass production (Edwards et al. 2006). Thus, it can be expected that P addition increases legume biomass production and symbiotic N_2 fixation. Furthermore, it has been observed that simultaneous additions of N and P (NP) can offset the negative effect of N addition on N_2 fixation resulting in higher symbiotic N_2 fixation compared to only N addition in tropical leguminous trees (Zheng et al. 2016). However, a recent study revealed that P addition enhances legume abundance in grasslands but does not mitigate the negative N effect when both elements are added simultaneously (Tognetti et al. 2021). Nevertheless, the extent to which the NP addition affects the N_2 fixation rates in grasslands remains to be studied.

Most studies on the effect of N and P addition on N_2 fixation in grasslands differ in experimental design, element addition rate, type of fertilizer used, and sampling procedure which leads to biases and uncertainties hampering our understanding of the main drivers of symbiotic N_2 fixation at the global scale (Zheng et al. 2019). Therefore, a standardized and globally replicated experiment is needed to gain insight into N_2 fixation. Recent work using a standardized and globally replicated experiment has shown that legume biomass production declines with N

addition (Tognetti et al. 2021), but the extent to which N and P additions affect symbiotic N_2 fixation rates was not evaluated.

Here, we use a standardized evaluation of symbiotic N_2 fixation in a globally coordinated grassland experiment, replicated at sites spanning a wide range of climatic and edaphic characteristics, to shed light on the response of symbiotic N_2 fixation in grasslands to N and P inputs. To evaluate the influence of single and combined N and P additions under various environmental conditions, we studied symbiotic N_2 fixation in 17 natural and semi-natural grassland sites with natural abundance of legumes (i.e. legumes were not deliberately introduced for the study) located on four continents. We determined symbiotic N_2 fixation based on the natural abundance of ^{15}N in shoot plant biomass. This approach relies on the distinct isotopic N signature of atmospheric N_2 and reactive soil N, which affects the plant N isotopic signature, depending on the source from which plants take up N (Amarger et al. 1979; Hoegberg 1997). We hypothesized that (i) N addition leads to a reduction in symbiotic N_2 fixation since it decreases legume biomass and N_2 fixation per unit legume biomass (i.e., the content of N derived from the atmosphere per unit of legume biomass), (ii) P addition enhances symbiotic N_2 fixation by increasing legume biomass and N_2 fixation per unit legume biomass, and (iii) the combined application of N and P increases N_2 fixation by offsetting the N-induced P deficiency caused by N application.

Material and methods

Study sites

The 17 study sites (Table 1, Figure S1) are experimental grasslands part of the Nutrient Network Global Research Cooperative (NutNet, <https://nutnet.org>) (Borer et al. 2014, 2017) and were selected according to the criterion that legumes were recorded in a minimum of six out of the 12 experimental plots (Table S1). The study sites are natural and semi-natural grasslands representing the regional flora with natural abundance of legumes (i.e. legumes were not deliberately introduced for the study). The selected sites are distributed across four continents (Table 1, Figure S1) covering a wide range of climatic conditions: Mean annual

Table 1 Continent, country, region/state, site name, latitude and longitude, elevation, mean annual precipitation (MAP), mean annual temperature (MAT), potential evapotranspiration (PET), estimated atmospheric nitrogen deposition (Ndep), year of study establishment and seasons of element addition (i.e., seasons from the establishment to the sample collection) of the 17 NutNet sites included in the study

Continent	Country	Region/State	Site	Latitude (°)	Longitude (°)	Elevation (m)	MAP (mm)	MAT (°C)	PET (mm)	Ndep (kg N ha ⁻¹)	Year of establishment	Seasons of element addition
Africa	South Africa	KwaZulu-Natal	Mt. Gilboa	29.3 S	30.3 E	1748	867	13.1	1194	5.6	2010	10
Africa	South Africa	KwaZulu-Natal	Ukulinga	29.7 S	30.4 E	843	838	18.1	1393	5.6	2009	7
America	Argentina	Santa Cruz	Potrok Aike	51.0 S	70.4 W	150	243	6.3	2923	0.5	2015	2
America	Canada	Ontario	Koffler Reserve	44.0 N	79.5 W	301	834	6.4	835	10.0	2010	5
America	USA	California	Hopland REC	39.0 N	123.1 W	598	939	12.3	1194	2.5	2007	11
America	USA	Kentucky	Spindletop	38.1 N	84.5 W	271	1166	12.5	1139	10.7	2007	10
America	USA	Oregon	Bunchgrass	44.3 N	122.0 W	1318	2160	5.5	860	3.2	2007	11
America	USA	Texas	Temple	31.0 N	97.3 W	184	870	19.1	1463	7.5	2007	8
Europe	Finland	Lapland	Saana	69.0 N	20.8 E	600	400	-3.1	339	3.5	2014	3
Europe	Germany	Bavaria	Bayreuth	49.9 N	11.6 E	340	724	8.3	756	14.6	2016	3
Europe	Germany	Saxony-Anhalt	Bad Lauchstädt	51.4 N	11.9 E	120	489	8.9	117	14.8	2015	2
Europe	Germany	Thuringia	Jena	50.9 N	11.5 E	320	597	8.0	724	14.6	2013	4
Europe	Portugal	Ribatejo	Companhia das Lezírias	38.0 N	8.0 W	200	642	16.5	1220	3.0	2012	4
Europe	Switzerland	Graubünden	Val Mustair	46.6 N	10.4 E	2320	950	0.3	442	21.7	2008	7
Europe	United Kingdom	North West	Lancaster	54.0 N	2.6 W	180	1222	8.0	599	10.2	2008	9
Oceania	Australia	Northern Territory	Kidmand Springs	16.1 S	131 E	87	749	27.3	2046	1.9	2014	1
Oceania	Australia	Western Australia	Pingelly Pad-dock	32.5 S	117.0 E	338	446	16.2	1427	1.0	2013	2

temperature (MAT) ranged between -3.1 and 27.3 °C and the mean annual precipitation (MAP) from 243 to 1222 mm. Sites were located between 87 and 2320 m above sea level (Table 1).

An identical experiment design is replicated at each site with four treatments: control (Ctrl; no element addition), N addition (N; 100 kg N ha⁻¹ yr⁻¹ as slow-release urea with $\delta^{15}\text{N}$ close to 0‰ (Choi et al. 2017)), P addition (P; 100 kg P ha⁻¹ yr⁻¹ as triple superphosphate (Ca(H₂PO₄)₂·H₂O)), and combined N and P addition (NP) (100 kg ha⁻¹ yr⁻¹ of both N and P). All treatments are replicated three times (n=3) at each site, and the experiments are organized in a randomized block design with 25 m² plots (5×5 m). All sites follow the same experimental design and sampling protocol. Further details about the experimental design and sampling can be found in Borer et al. (2014)

The climatic data were derived from Hijmans et al. (2005) based on the location of each site (Table 1). Nitrogen deposition (as kg N ha⁻¹ yr⁻¹) was estimated based on the location of each site (longitude and latitude) using the model output of Ackerman et al. (2019) for the year 2016 (Table 1). The soil properties of the control treatment at the time of establishment

of the experiment are summarized in the Table 2. The methods used for soil analysis have been described in Seabloom et al. (2021).

Plant sampling

Aboveground biomass was sampled annually at the time of peak biomass using a standardized protocol (Borer et al. 2014). For the present study, we used the aboveground biomass of a single sampling between the years 2015 and 2020 depending on the site (as detailed in Table 1). Two 10×100 cm strips (covering area of 0.2 m²) of vegetation were clipped directly above the soil surface in a subplot of 1×1 m within each plot. The clipped plant biomass was sorted into the three functional groups: grasses, non-leguminous forbs and legumes, and oven-dried at 60 °C to a constant mass prior to weighing. Hereafter, we will refer to aboveground plant biomass as plant biomass. Representative subsamples of the biomass of the three plant functional groups from all plots were sent to the University of Bayreuth (Germany) for further analyses. The two most abundant grass, forb and legume species based on the cover estimates in the control

Table 2 Selected soil properties (0–10 cm) at the time of experiment establishment of the 17 Nutrient Network (NutNet) sites evaluated in the study

Continent	Country	Site	TOC (g kg ⁻¹)	TN	C:N ratio	pH	TP (mg kg ⁻¹)	Ca	Mg	K	Na
Africa	South Africa	Mt. Gilboa	20.6	1.15	17.8	5.06	16.6	227	43.1	126	36.2
Africa	South Africa	Ukulinga	47.2	3.29	14.4	5.40	38.9	2028	710	178	65.4
America	Argentina	Potrok Aike	38.9	2.87	13.6	6.46	23.4	1862	476	292	76.6
America	Canada	Koffler Reserve	19.5	1.53	12.8	7.41	13.8	4018	132	50.6	52.8
America	USA	Hopland REC	24.7	2.00	12.2	6.63	26.0	1426	297	187	23.0
America	USA	Spindletop	26.5	2.56	10.3	6.40	233	2469	241	77.8	30.4
America	USA	Bunchgrass	88.7	6.10	14.5	5.54	13.8	291	39.9	94.2	28.8
America	USA	Temple	98.3	3.69	26.7	7.69	20.7	19,005	220	437	32.2
Europe	Finland	Saana	119	4.88	23.4	6.53	23.4	4026	813	54.9	29.8
Europe	Germany	Bayreuth	14.9	1.20	12.0	4.73	47.1	672	59.7	96.9	35.1
Europe	Germany	Bad Lauchstädt	35.2	1.84	19.2	7.13	44.9	2546	186	203	20.8
Europe	Germany	Jena	58.5	5.33	11.0	7.43	177	9268	292	1145	28.0
Europe	Portugal	Companhia das Lezírias	18.0	1.21	15.1	5.92	34.4	843	54.8	85.3	23.9
Europe	Switzerland	Val Mustair	77.1	6.05	12.8	5.49	48.0	1422	277	103	21.6
Europe	United Kingdom	Lancaster	20.5	1.10	18.5	4.78	34.3	1486	121	112	44.1
Oceania	Australia	Kidmand Springs	25.4	n.d	n.d	7.90	2.92	4774	1777	478	55.1
Oceania	Australia	Pingelly Paddock	20.1	1.28	15.9	6.00	14.9	1149	99.1	129	34.5

TOC, Total organic carbon; TN, Total nitrogen; C:N ratio, Carbon to nitrogen ratio (mass: mass); TP, Total phosphorus; Ca, Calcium; Mg, Magnesium; K, Potassium; Na, Sodium; n.d., Not determined

plots for each year of sampling at each site are shown in Table 3. The cover estimate of each vascular plant was estimated visually in a 1 m² subplot within each treatment plot when plant biomass was highest (peak biomass). In addition, the sites were classified as grassland with perennial or annual legumes according to the type (perennial or annual) of the two most abundant legume species (Table 3).

Plant C, N and P concentration and stable isotope determination ($\delta^{15}\text{N}$)

In total, 553 dried plant samples were processed (201 grasses, 185 forbs and 167 legumes, Table S1). Plant biomass of each functional group was cut with scissors, homogenized and ground in a ball mill. The total C and N concentration and the isotopic composition were analyzed using continuous-flow isotope ratio mass spectrometry (NA 1108 elemental Analyzer, CE Instruments, Milano, Italy) coupled via ConFlo III open-split interface (Finnigan MAT, Bremen, Germany) to a delta S isotope ratio mass spectrometer (Finnigan MAT, Bremen, Germany) at the University of Bayreuth. The isotopic composition of N was expressed in δ notation, which represents the ‰ of variation compared to the international standard for natural ¹⁵N abundance measurements (atmospheric N isotope ratio). In addition, the total P concentration of plant biomass was determined by ICP-OES (Vista-Pro radial, Varian, Aschaffenburg, Germany) after nitric acid digestion.

Calculations

The percentage of legume N in aboveground biomass derived from atmospheric N₂ fixation (%Ndfa), via N₂ fixation by legumes, was calculated following the approach described by Amarger et al. (1979) and Hoegberg (1997) using Eq. 1:

$$\text{Ndfa (\%)} = \frac{(\delta^{15}\text{N}_{\text{reference}} - \delta^{15}\text{N}_{\text{legume}})}{(\delta^{15}\text{N}_{\text{reference}} - \text{B})} \times 100 \quad (1)$$

where $\delta^{15}\text{N}_{\text{reference}}$ is the $\delta^{15}\text{N}$ of a non-fixing reference plant, $\delta^{15}\text{N}_{\text{legume}}$ is the $\delta^{15}\text{N}$ of the legume

aboveground biomass (called legume $\delta^{15}\text{N}$ hereafter) in the evaluated plot, and B is the legume $\delta^{15}\text{N}$ fully relying on atmospheric N₂ fixation which accounts for any internal isotopic fractionation of the legume (Hoegberg 1997). We considered the mean of the $\delta^{15}\text{N}$ of the two non-fixing functional groups (grasses and forbs) as the reference, which was calculated separately for each plot at each site (and is called reference $\delta^{15}\text{N}$ hereafter).

We used the lowest legume $\delta^{15}\text{N}$ value of all plots at each site as the B value (Eq. 1), similar to previous studies (Hansen and Vinther 2001; West et al. 2005; Roscher et al. 2011; Oberson et al. 2013). This approach relies on the assumption that the legumes in the plot with the lowest legume $\delta^{15}\text{N}$ receive 100% of the N from symbiotic N₂ fixation. We used the lowest legume $\delta^{15}\text{N}$ from all four treatments because the legume $\delta^{15}\text{N}$ was not significantly affected by element addition (see below in the Results section), similar to previous studies that considered different element addition treatments (Oberson et al. 2013; Tzanakakis et al. 2017).

We detected a relative ¹⁵N-depletion in reference $\delta^{15}\text{N}$ compared to legume $\delta^{15}\text{N}$ (i.e., lower reference $\delta^{15}\text{N}$ than legume $\delta^{15}\text{N}$) at seven sites [Koffler (Canada), Hopland, Spindletop and Bunchgrass (USA), Bad Lauchstädt and Bayreuth (Germany) and Val Mustair (Switzerland)] which challenged the estimation of %Ndfa using Eq. 1 because it resulted in negative %Ndfa estimates. We observed a decrease in reference $\delta^{15}\text{N}$ (mean of grasses and forbs) with increasing elevation of the study site ($r^2=0.301$, $p=0.024$, Figure S2), and no significant relationship between elevation and legume $\delta^{15}\text{N}$ ($p=0.337$). This observation is consistent with previous studies, showing that $\delta^{15}\text{N}$ of non-fixing plants decreases with increasing elevation (Jacot et al. 2000; Craine and Lee 2003; Huber et al. 2007; Zhou et al. 2016). We adjusted the reference $\delta^{15}\text{N}$ for the effect of elevation (elevation adj. $\delta^{15}\text{N}_{\text{reference}}$), assuming that all sites would be located at an elevation of 0 m a.s.l., using the slope of the regression line describing the relationship between the $\delta^{15}\text{N}$ of the reference and elevation (see Figure S2), as:

$$\text{Elevation adj. } \delta^{15}\text{N}_{\text{reference}} (\text{‰}) = \delta^{15}\text{N}_{\text{reference}} (\text{‰}) - [-0.002 * \text{Elevation(m)}] \quad (2)$$

Table 3 Names of the two most abundant grass, forb and legume species and the type of dominating legumes (annual or perennial) at each site according to the cover estimates in the three control plots of each site for the year of sampling. In brackets, mean values of cover estimates (%) of each species in the control plots is given

Continent	Country	Site	Grasses	Forbs	Legumes	Legume type*
Africa	South Africa	Mt. Gilboa	<i>Diheteropogon filifolius</i> (Nees) Clayton (20)	<i>Protea simplex</i> E. Phillips (7)	<i>Eriosema distinctum</i> N.E.Br (18)	Perennial
			<i>Heteropogon contortus</i> (L.) P.Beauv. (20)	<i>Hypoxis gerrardii</i> Baker (4)	<i>Pearsonia grandifolia</i> (Bolus) Polhill (2)	Perennial
Africa	South Africa	Ukulinga	<i>Eragrostis curvula</i> (Schrad.) Nees (33)	<i>Scabiosa columbaria</i> L. (12)	<i>Eriosema cordatum</i> E. May. (4)	Perennial
			<i>Tristachya leucothrix</i> Trin. ex Nee (32)	<i>Helichrysum pilosellum</i> (L. f.) Less. (7)	<i>Rhynchosia minima</i> (L.) DC. (1)	Perennial
America	Argentina	Potrok Aike	<i>Stipa speciosa</i> Trin. & Rupr. (23)	-	<i>Adesmia lotoides</i> Hook. f. (<1)	Perennial
			<i>Poa spiciformis</i> (Steud.) (8)	-	<i>Lathyrus nervosus</i> Lam. (<1)	Perennial
America	Canada	Koffler Reserve	<i>Bromus inermis</i> Leyss. (17)	<i>Asclepias syriaca</i> L. (3)	<i>Vicia tenuifolia</i> Roth. (2)	Perennial
			<i>Poa spp.</i> (15)	<i>Euthamia graminifolia</i> Nutt. Ex Cass. (2)	-	-
America	USA	Hopland REC	<i>Briza máxima</i> L. (72)	<i>Carduus pycnocephalus</i> L. (3)	<i>Lupinus nanus</i> Douglas ex Benth. (3)	Annual
			<i>Elymus glaucus</i> Buckley (7)	<i>Sisyrinchium bellum</i> S.Watson (2)	<i>Trifolium hirtum</i> All (3)	Annual
America	USA	Spindletop	<i>Dactylis glomerata</i> L. (35)	<i>Erigeron annuus</i> L. (8)	<i>Vicia grandiflora</i> Scop. (12)	Annual
			<i>Festuca arundinacea</i> Schreb. (9)	<i>Plantago lanceolata</i> L. (5)	<i>Trifolium pratense</i> L. (5)	Perennial
America	USA	Bunchgrass	<i>Carex pensylvanica</i> Lam. (36)	<i>Phlox difusa</i> Benth. (40)	<i>Lupinus latifolius</i> Lindl. Ex J. Agardh (27)	Perennial
			<i>Poa pratensis</i> L. (7)	<i>Penstemon procerus</i> Dougl. ex Graham (10)	-	-
America	USA	Temple	<i>Ambrosia trifida</i> L. (38)	<i>Schizachyrium scoparium</i> (Michx.) Nash (15)	<i>Psoralidium tenuiflorum</i> (Pursh) Rydb. (5)	Perennial
			<i>Stenaria nigricans</i> (Lam.) (17)	<i>Sorghum halepense</i> (L.) Pers.(10)	<i>Mimosa nuttallii</i> (DC.) B.L. Turner (3)	Perennial
Europe	Finland	Saana	<i>Elymus mutabilis</i> (Drobow) Tzvelev (3)	<i>Geranium sylvaticum</i> L. (70)	<i>Astragalus alpinus</i> L. (<1)	Perennial
			<i>Melica nutans</i> L. (2)	<i>Trollius europaeus</i> L. (20)	-	-
Europe	Germany	Bayreuth	<i>Festuca rubra</i> L. (9)	<i>Achillea millefolium</i> L. (3)	<i>Trifolium dubium</i> SIBTH. (3)	Annual
			<i>Luzula campestris</i> (L.) DC. (3)	<i>Scorzoneroidea autumnalis</i> (L.) Moench (2)	<i>Vicia hirsuta</i> (L.) Gray (2)	Annual
Europe	Germany	Bad Lauchstädt	<i>Vulpia myuros</i> (L.) C.C. Gmel. (12)	<i>Picris hieracioides</i> L. (33)	<i>Medicago lupulina</i> L. (23)	Annual
			-	<i>Taraxacum officinale</i> F.H.Wigg. (3)	<i>Trifolium dubium</i> SIBTH. (3)	Annual

Table 3 (continued)

Continent	Country	Site	Grasses	Forbs	Legumes	Legume type*
Europe	Germany	Jena	<i>Lolium perenne</i> L. (21)	<i>Taraxacum officinale</i> F.H.Wigg. (22)	<i>Trifolium dubium</i> SIBTH. (30)	Annual
			<i>Poa pratensis</i> L. (11)	<i>Crepis biennis</i> L. (22)	<i>Medicago lupulina</i> L. (<1)	Annual
Europe	Portugal	Companhia das Lezírias	<i>Agrostis pourretii</i> WILLD. (9)	<i>Tolpis barbata</i> (L.) GAERTN. (26)	<i>Ornithopus compressus</i> L. (13)	Annual
			<i>Vulpia geniculata</i> (L.) Link (4)	<i>Tuberaria guttata</i> (L.) Fourr. (13)	<i>Trifolium arvense</i> L. (4)	Annual
Europe	Switzerland	Val Mustair	<i>Agrostis alpina</i> Scop. (9)	<i>Hieracium pilosella</i> L. (11)	<i>Trifolium pratense</i> L. (10)	Perennial
			<i>Festuca halleri</i> All. (5)	<i>Carlina acaulis</i> L. (5)	<i>Lotus alpinus</i> (Ser.) Schleich. ex Ramond (7)	Perennial
Europe	United Kingdom	Lancaster	<i>Agrostis capillaris</i> L. (47)	<i>Ranunculus repens</i> L. (4)	<i>Trifolium repens</i> L. (5)	Perennial
			<i>Holcus lanatus</i> L. (34)	<i>Rumex acetosa</i> L. (1)	-	-
Oceania	Australia	Kidmand Springs	<i>Chrysopogon fallax</i> S.T.Blake (35)	<i>Polymeria ambigua</i> R.Br. (1)	<i>Rhynchosia minima</i> (L.) DC. (<1)	Perennial
			<i>Panicum decompositum</i> R.Br. (35)	<i>Trichodesma zeylanicum</i> (BURM.F.) R.Br. (<1)	<i>Neptunia spp.</i> (<1)	Perennial
Oceania	Australia	Pingelly Pad-dock	<i>Avena barbata</i> POTT EX. LINK (17)	<i>Erodium botrys</i> (CAV.) BERTOL. (13)	<i>Trifolium subterraneum</i> L. (20)	Annual
			<i>Vulpia myuros</i> (L.) C.C.Gmel. (11)	<i>Hypochaeris glabra</i> L. (13)	<i>Trifolium arvense</i> L. (<1)	Annual

*The two most abundant legume species were of the same type at each site (i.e. they were either both annual or perennial) except in Spindletop (USA) where the most abundant (*Vicia grandiflora*) was annual and the second most abundant (*Trifolium pratense*) was perennial. Because *Vicia grandiflora* was more abundant and, unlike *Trifolium pratense*, was present in all three blocks, the site was considered as annual

Next, Ndfa (%) was calculated based on the Elevation adj. $\delta^{15}\text{N}_{\text{reference}}$ as follows:

$$\text{Ndfa (\%)} = \frac{(\text{Elevation adj. } \delta^{15}\text{N}_{\text{reference}} - \delta^{15}\text{N}_{\text{legume}})}{(\text{Elevation adj. } \delta^{15}\text{N}_{\text{reference}} - \text{B})} \times 100 \quad (3)$$

Further details about the elevation adjustment and %Ndfa calculation are presented as Supporting Information.

Legume N uptake was calculated based on the legume N concentration at peak biomass as:

$$\begin{aligned} \text{Legume N uptake (mg N g}^{-1} \text{ yr}^{-1}) \\ = \text{legume N concentration (mg N g}^{-1}) \times \text{yr}^{-1} \end{aligned} \quad (4)$$

N_2 fixation per legume biomass was calculated for the plots with legumes as:

$$\begin{aligned} \text{N}_2 \text{ fixation per unit legume biomass (mg N g}^{-1} \text{ yr}^{-1}) \\ = \text{Legume N uptake (mg N g}^{-1} \text{ yr}^{-1}) \\ \times \text{Ndfa (\%)} \times 0.01 \end{aligned} \quad (5)$$

Symbiotic N_2 fixation per area grassland was calculated as follows:

$$\begin{aligned} \text{N}_2 \text{ fixation per area (g N m}^{-2} \text{ yr}^{-1}) \\ = \text{legume biomass (g m}^{-2}) \\ \times \text{N}_2 \text{ fixation per legume biomass} \\ \text{(mg N g}^{-1} \text{ yr}^{-1}) \end{aligned} \quad (6)$$

Symbiotic N₂ fixation per total grassland biomass (with grassland biomass being the sum of the biomasses of all three functional groups) was calculated for all plots as:

$$\begin{aligned} \text{N}_2 \text{ fixation per total biomass (mg N g}^{-1}\text{yr}^{-1}) \\ = \text{N}_2 \text{ fixation per area (g N m}^{-2}\text{yr}^{-1}) \\ \times 1000 / \text{total biomass (g m}^{-2}) \end{aligned} \quad (7)$$

We calculated means and standard errors of all four treatments across all 17 sites. If legumes were absent in a plot, the legume N stock (in g ha⁻¹) or legume N uptake of this plot was assumed to be zero. If the legume biomass was zero, we assumed that symbiotic N₂ fixation per unit biomass or area was also zero. When calculating the mean of the N or P concentration of the biomass of all three functional groups, we considered only plots with biomass of the respective functional group. Similarly, when calculating the mean of symbiotic N₂ fixation per legume biomass, only plots with legume biomass > 0 and valid %Ndfa were considered. In contrast, when calculating symbiotic N₂ fixation per unit area of grassland or total grassland biomass, all plots with legumes and valid %Ndfa as well as plots without legumes were considered.

Statistical analyses

Data were analyzed using linear mixed models with the software SPSS 27 (IBM SPSS, Inc., Chicago, USA). Because data were not normally distributed (Shapiro–Wilk-test, $p > 0.05$), all variables except the $\delta^{15}\text{N}$ values (including both positive and negative values) were log-transformed. The different element addition treatments (Ctrl, N, P, NP) and site- and plot-level covariates were used as fixed factors, and block as a random factor where block was nested within site. The site-level covariates included in the linear mixed model were MAP, MAT, water availability index (MAP/potential evapotranspiration), the estimates of N deposition and the legume proportion of biomass in the control treatments at each site. Because sites were set up in different years, the number of years of element addition was considered as a site-level covariate (redundant in most of the evaluated parameters with no significant effect). Soil properties summarized in Table 2 (except total organic carbon (TOC), which

was highly correlated with total nitrogen (TN)) were included in the linear mixed model as plot-level covariates. The interactions between treatment and the plot-level covariates were initially considered in the model, although after a selection based on Akaike Information Criterion only the interactions ‘treatment x TN’ and ‘treatment x soil pH’ remained in the final model as covariates. When a significant treatment effect ($p < 0.05$) was found, LSD post hoc test ($p < 0.05$) was used for comparison of means of the element addition treatments.

Additionally, we evaluated how the type of dominating legumes (perennial or annual) at each site affected legume biomass, legume N concentration, legume $\delta^{15}\text{N}$, %Ndfa and symbiotic N₂ fixation per area of grassland. A linear mixed model was used with treatment (Ctrl, N, P and NP), type of dominating legumes (annual or perennial), their interaction and the site- and plot-level covariates as fixed factors, and block as a random factor nested within site. The model was performed as previously described.

We calculated the response to nutrient addition of legume biomass, %Ndfa and N₂ fixation per unit area as:

$$\text{Response} = \text{Ln} ((Y_{\text{treatment}} + 1)/(Y_{\text{control}} + 1)) \quad (8)$$

where $Y_{\text{treatment}}$ is the value of legume biomass, %Ndfa or N₂ fixation per unit area in the N, P or NP addition treatment and Y_{control} is the mean value of legume biomass, %Ndfa or N₂ fixation per unit area in the control. The response was calculated separately for each site. We added 1 to the numerator and denominator to remove zeros before the logarithmic transformation.

We performed stepwise multiple regression analyses to evaluate the impact of site- and plot-level covariates on the response of legume biomass, %Ndfa and N₂ fixation per unit area to nutrient addition. Stepwise multiple regressions analyses were performed using the site-scale factors (MAP, MAT, water availability, N deposition and the legume proportion of biomass in the control treatment at each site) and plot-scale soil properties (TN, carbon-to-nitrogen ratio (C:N ratio), available P, and soil pH). Collinearity was evaluated based on the variance inflation factor. The multiple regression analyses were performed separately for the three different element addition treatments (i.e., N, P and NP).

Results

Plant aboveground biomass

On average across sites, N and P addition increased total plant biomass (the sum of the biomasses of all three functional groups) by 32 and 28%, respectively, compared to the control (Fig. 1). The combined addition of N and P increased the total plant biomass by 72%, from a mean of 3040 kg ha⁻¹ in the control to 5222 kg ha⁻¹ in the NP treatment. The biomass of both the grasses and the forbs increased significantly with NP addition (Fig. 1) but did not respond significantly to N or P addition alone.

Legume biomass was highly variable among sites and ranged from 0.1 kg ha⁻¹ (Potrock, Argentina) to 1082 kg ha⁻¹ (Bad Lauchstädt, Germany) in the control treatment (Table S2). Compared to the control, biomass of legumes was reduced by 65% and 45% in the N and NP treatments, respectively, while in the P treatment was increased by 77% (although this difference was not statistically significant) (Fig. 1). Compared to the P treatment, biomass of legumes was significantly reduced by 81% and 69% by N and NP

addition. The percent of legumes in the total plant biomass (the sum of the biomasses of all three functional groups) was 9.9% in the control, 4.7% in the N, 12.0% in the P, and 3.3% in the NP treatment (Fig. 1). No significant interaction between the treatments and the type of dominant legumes (annual or perennial) at each site was observed on legume biomass, although legume biomass was significantly higher in the grasslands dominated by annual than by perennial legumes (Figure S3A). Similar differences between P and N and NP treatments were observed in the response ratio to nutrient addition of legume biomass (Figure S4A).

Nitrogen and phosphorus concentrations in aboveground biomass

The addition of N significantly increased plant N concentrations (Fig. 2A), while the addition of P significantly increased plant P concentrations (Fig. 2B) in both grasses and forbs compared to the control. In contrast, legume N concentration was not affected by N addition, whereas legume P concentration was




Fig. 1 Aboveground biomass at peak biomass of grasses, forbs, and legumes as well as total biomass (grasses, forbs and legumes together) as affected by single and combined N and P addition. Bars show means ($n=51$) with standard errors. Ctrl, control; N, nitrogen addition; P, phosphorus addition; NP,

nitrogen and phosphorus addition. All plots were included in this calculation irrespective of amount of biomass of the functional groups. Different letters indicate significant differences ($p < 0.05$) among treatments, tested separately for each plant functional group

enhanced by both the P and NP treatments (Fig. 2A, B). No significant interaction between the treatments and the type of dominating legumes (annual or perennial) was observed in legume N concentration (Figure S3B). The biomass N:P ratio in the control treatment was 8.5 for grasses, 7.3 for forbs, and 13.9 for legumes (Fig. 2C). The N:P ratio of legumes was significantly reduced by the addition of P (N:P=8.7) and NP (N:P=7.9) compared to the control and the N treatment (Fig. 2C). Similarly, P addition decreased the N:P ratio of grasses and forbs (Fig. 2C).

The legume N stock was significantly higher in the control (0.71 g N m^{-2}) and P treatment (1.34 g N m^{-2}) than in the N (0.24 g N m^{-2}) and NP (0.36 g N m^{-2}) treatment (Figure S5A). The P stock of grasses and forbs was increased by P and NP addition in comparison to the control, while the P stock of legumes was only increased in the P treatment compared to the control (Figure S5B).

Plant isotopic composition

We observed a decrease in $\delta^{15}\text{N}$ of the reference functional groups (grasses and forbs) in the control treatment with increasing elevation of the study site ($r^2=0.301$, $p=0.024$, Figure S2). Therefore, we adjusted the reference $\delta^{15}\text{N}$ of all plots and treatments for elevation (see Sect. Calculations). After the recalculation of the reference $\delta^{15}\text{N}$, the mean of the elevation-adjusted $\delta^{15}\text{N}_{\text{reference}}$ was $+0.81\text{‰}$ in the control treatment (Table S3). The $\delta^{15}\text{N}_{\text{reference}}$ before and after the elevation-adjustment was significantly higher in the N and NP treatments than in the control and P treatment (Table S3). The legume $\delta^{15}\text{N}$ across all sites was unaffected by treatments (Table S3).

Symbiotic N_2 fixation per unit legume biomass

The mean %Ndfa in the control treatment was 65.8% across all 17 sites (Fig. 3A). No significant difference in %Ndfa among treatments was found. However, %Ndfa was slightly higher in the P and NP treatments (69.8 and 70.9%, respectively) than in the control and N treatments (65.8 and 64.2%, respectively). Similarly, no significant difference was observed in the %Ndfa response to nutrient addition (Figure S4B). The response of %Ndfa to NP addition was positively related to the proportion of legumes in the total




Fig. 2 Plant aboveground nitrogen (A) and phosphorus (B) concentrations and plant aboveground nitrogen to phosphorus ratio (mass: mass) (C) of grasses, forbs and legumes as affected by single and combined N and P addition. Bars show means with standard errors. White numbers at the base of each column indicate the number of replicates. Only plots with biomass of the functional group >0 were included in the calculation of the element concentrations of each functional group. Ctrl, control; N, nitrogen addition; P, phosphorus addition; NP, nitrogen and phosphorus addition. Different letters indicate significant differences ($p < 0.05$) among treatments, tested separately for each plant functional group




Fig. 3 Percentage of legume biomass-nitrogen derived from atmospheric N₂ fixation (%Ndfa) (A), symbiotic N₂ fixation by legumes per legume biomass (B) as well as symbiotic N₂ fixation by legumes per unit area of grassland (C) and per total grassland biomass (D) as affected by single and combined N and P addition. White number at the base of each bar shows the number of replicates. Only plots with legume biomass > 0

were included in panels A and B, whereas panels C and D also include plots with no legume biomass. Bars represent means with standard errors. Ctrl, control; N, nitrogen addition; P, phosphorus addition; NP, nitrogen and phosphorus addition. Treatments with different letters represent significant differences at $p < 0.05$

biomass in the control treatment, initial soil N and water availability index, while it was negatively related to N deposition (Table 4, Table S4). There were no significant linear regression models ($p > 0.05$) for the %Ndfa response to N or P addition. We observed a significantly higher %Ndfa in the ten sites dominated by perennial legumes (74.0%) compared to the seven sites dominated by annual legumes (58.0%) (Fig. 4). In addition, the interaction between treatment and type of dominating legumes revealed that the single N addition limited the differences in %Ndfa between annual and perennial legumes (Fig. 4).

Mean N₂ fixation per unit legume biomass in the control treatment across all 17 study sites was

18.1 mg N g⁻¹ yr⁻¹ legume biomass (Fig. 3B). No significant difference among treatments in N₂ fixation per legume biomass was observed due to the lack of element addition effect on legume N concentration and %Ndfa. However, N₂ fixation per unit legume biomass was slightly higher in the NP treatment (20.2 mg N g⁻¹ legume biomass yr⁻¹) than in the other three treatments (Fig. 3B).

Symbiotic N₂ fixation per unit area grassland

Symbiotic N₂ fixation per unit area in the control treatment across all 17 study sites was

Table 4 Regression models of the response of legume biomass, the proportion of N derived from atmosphere (%Ndfa), and symbiotic N₂ fixation per unit area to element addition (nitrogen (N), phosphorus (P) and their combined application (NP)) as a function of site-scale environmental factors (MAP, MAT, water availability (Aw), N deposition (Ndep) and leg-

ume proportion of biomass in the control treatment at each site (prop)) and plot-scale soil properties (total nitrogen (TN), soil carbon to nitrogen ratio (C:N), available phosphorus (P) and soil pH). A dash (-) indicates that no significant ($p < 0.05$) model was found

Dependent variable (y)	Response to	Regression model	Adjusted r^2	p -value
Response of legume biomass	N addition	$y = 5.792 - 0.752\text{pH} - 0.128\text{MAT} - 0.110\text{Ndep} - 2.150\text{TN} + 0.006\text{P}$	0.660	0.001
	P addition	$y = 2.360 - 6.091\text{TN} - 0.082\text{MAT}$	0.431	0.001
	NP addition	$y = 1.570 - 6.009\text{TN} - 0.111\text{MAT}$	0.472	0.001
Response of %Ndfa	N addition	-	-	-
	P addition	-	-	-
	NP addition	$y = -0.061 + 0.007\text{Prop} - 0.027\text{Ndep} + 0.449\text{TN} + 0.073\text{Aw}$	0.613	0.001
Response of N ₂ fixation	N addition	$y = 3.389 - 0.481\text{pH} - 0.048\text{Ndep} - 0.036\text{MAT} - 0.001\text{MAP}$	0.550	0.001
	P addition	$y = 0.529 - 2.430\text{TN}$	0.186	0.009
	NP addition	$y = 0.817 - 2.403\text{TN} - 0.042\text{MAT} - 0.046\text{Ndep}$	0.406	0.001




Fig. 4 Percentage of legume biomass-nitrogen derived from atmospheric N₂ fixation (%Ndfa) as affected by single and combined N and P addition (Treatment), the type of dominating legumes (annual or perennial) (Type) and their interaction (T x T). Bars show means with standard errors. Numbers at the base of each column indicate the number of replicates.

Ctrl, control; N, nitrogen addition; P, phosphorus addition; NP, nitrogen and phosphorus addition. * and *** indicate significant effect at $p < 0.05$ and $p < 0.001$, respectively; n.s. not significant. Different letters indicate significant differences ($p < 0.05$) between annual and perennial within each treatment

3.5 kg N ha⁻¹ yr⁻¹ (Fig. 3C). It ranged from 0.002 kg ha⁻¹ yr⁻¹ (Potrock, Argentina) to 11.9 kg N ha⁻¹ yr⁻¹ (Bad Lauchstädt, Germany). Across all sites, N₂ fixation per area

was 1.39 kg N ha⁻¹ yr⁻¹ in the N treatment and 2.13 kg N ha⁻¹ yr⁻¹ in the NP treatment. Thus, N₂ fixation was significantly reduced in the N and NP treatment by 60 and 39%, respectively,

compared to the control. Similarly, N_2 fixation per unit area was also 63% lower in the N and 43% lower in the NP treatment than in the P treatment ($3.71 \text{ kg N ha}^{-1} \text{ yr}^{-1}$) (Fig. 3C). In contrast, P addition had no significant effect on N_2 fixation per area compared to the control. The number of years of element addition was a redundant site-level covariate in the linear mixed model, indicating that the different ages of the sites did not influence the estimation of N_2 fixation in the present study. In addition, no significant effect of the type of dominating legumes (annual or perennial) on N_2 fixation per area was observed (Figure S3D). Similarly, the response to N and NP addition of the N_2 fixation per unit area was significantly lower than the response to P addition (Figure S4C). Soil pH, N deposition, MAT and MAP were negatively related with the response of N_2 fixation per unit area to N addition ($r^2=0.550$, $p=0.001$, Table 4, Table S4). In addition, the response of N_2 fixation per unit area to P addition was negatively related with soil N ($r^2=0.162$, $p=0.009$, Table 4, Table S4), while the response of N_2 fixation per unit area to NP addition was positively related with soil N, MAT and N deposition ($r^2=0.406$, $p=0.001$, Table 4, Table S4).

The mean N_2 fixation per unit total biomass in the control treatment was $1.23 \text{ mg N g}^{-1} \text{ yr}^{-1}$ across all 17 study sites (Fig. 3D). The N_2 fixation per total biomass was reduced by 60% under N and 73% under NP addition compared to the control (Fig. 3D). Similarly, N_2 fixation per total biomass was lower in the N and NP treatments compared to the P treatment.

Discussion

Our results reveal that the addition of N decreased the rate of N_2 fixation per grassland area compared to the control, and this effect was not reversed by additional P amendment (Fig. 3C). The reduced N_2 fixation per area grassland was due to the reduction in legume biomass, and not an altered N_2 fixation rate per unit legume biomass. Tognetti et al. (2021) recently showed that legume biomass was negatively affected by N addition in grasslands on several continents. Our study goes further, demonstrating that this reduction in legume biomass causes the N_2 fixation rate per unit area to decrease from $3.50 \text{ kg N ha}^{-1} \text{ yr}^{-1}$

to $1.39 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ due to N addition across all sites. Further, we found that N_2 fixation per legume biomass was not significantly affected by N or P addition, which is an important finding since it suggests that the expression of the enzyme that fixes atmospheric N_2 is not downregulated in legumes in response to high availability of reactive N in the soil as observed by Menge and Hedin (2009).

$\delta^{15}\text{N}$ patterns in plant functional groups

We observed a significant negative correlation between site elevation and the $\delta^{15}\text{N}$ of the reference plants (grass and forbs; Figure S2), which suggests low soil $\delta^{15}\text{N}$ at high elevations. Similar observations were described in previous global reviews (Amundson et al. 2003) and in studies along altitudinal gradients (Vitousek et al. 1989; Jacot et al. 2000; Craine and Lee 2003; Huber et al. 2007; Zhou et al. 2016). The underlying reason is the relationship between elevation and MAT and MAP, two of the main drivers of plant $\delta^{15}\text{N}$ as described by Craine et al. (2009) and Zhou et al. (2016). The elevation-dependent $\delta^{15}\text{N}$ of non-fixer plants (reference plants) likely caused difficulties when applying the ^{15}N natural abundance method to determine symbiotic N_2 fixation about altitudinal gradients in previous studies (Vitousek et al. 1989; Jacot et al. 2000). The unique global design of this study allowed us to correct this elevation effect on reference plants and to determine symbiotic N_2 fixation across a large number of sites on different continents. The relationship between elevation and $\delta^{15}\text{N}$ of grasses and forbs identified here will likely be of use also in future studies. However, external inputs of ^{15}N -depleted N, such as cattle urine, large inputs of legume-derived N or atmospheric N deposition cannot be dismissed as other factors affecting $\delta^{15}\text{N}$ of grasses and forbs (Jacot et al. 2000; Hansen and Vinther 2001; Gehring and Vlek 2004). Further details about the elevation adjustment and ^{15}N natural abundance method are presented as Supporting Information.

Symbiotic N_2 fixation per area decreased by N addition

We found that N addition significantly reduced the rate of symbiotic N_2 fixation per area of grassland due to a reduction in legume biomass without

affecting the N_2 fixation rate per unit legume biomass. The negative effect of N addition on symbiotic N_2 fixation is consistent with previous single-site studies observing a reduction in symbiotic N_2 fixation by N addition (West et al. 2005; Nyfeler et al. 2011; Burchill et al. 2014; Tzanakakis et al. 2017) and a recent study showing that also non-symbiotic N_2 fixation in the soil is decreased by N addition at sites of the Nutrient Network experiment (Schleuss et al. 2021). Our results suggest that continuous anthropogenic N enrichment of grasslands can lead to a decrease in legume biomass production, which in the long-term can limit symbiotic N_2 fixation. The most plausible explanation for reduced rates of N_2 fixation per unit area due to N addition is that higher soil N availability allows grasses and forbs to outcompete legumes (via competition for light) and reduce legume biomass (Suding et al. 2005; Soussana and Tallec 2010; Tognetti et al. 2021). The size of the response of symbiotic N_2 fixation per area to N addition was affected by several abiotic factors. Addition of N reduced N_2 fixation per area more strongly at sites with higher soil pH, atmospheric N deposition, MAT, and MAP (Table 4, Table S4). The reasons for this could be that i) legumes and rhizobium strains from neutral and alkaline sites are less tolerant to soil acidification caused by urea addition (Hungria and Vargas 2000), that ii) N addition has a larger effect at sites where the anthropogenic N input through atmospheric N deposition is already large, and iii) N_2 fixation is more sensitive to N addition at sites where the N_2 fixation is not constrained by temperature or water availability (Houlton et al. 2008; Tognetti et al. 2021).

In contrast to our second hypothesis, we observed similar rates of symbiotic N_2 fixation per area of grassland in the control and P treatment, which can be attributed to a lack of P limitation of N_2 fixation in the control treatment as further indicated by the low N:P ratio of legume biomass (13.9) (Güsewell 2004). The reason for the lack of effect to P addition is likely that legumes have evolved very effective mechanisms to increase their P uptake from different soil P pools through the release of phosphatases or organic acids into the rhizosphere (Nuruzzaman et al. 2006). The negative relationship between the response of N_2 fixation per area and soil N to P addition (Table 4, Table S4) might indicate that at N-limited sites, legumes could invest the added P

in symbiotic N_2 fixation to overcome the N limitation (McKey 1994; Houlton et al. 2008; Soussana and Tallec 2010).

The application of P in combination with N did not counterbalance the negative impact of N addition on symbiotic N_2 fixation per area, in contrast to our third hypothesis. Our results indicate that the negative impact of N addition on symbiotic N_2 fixation per area is not a result of N-driven P deficiency because in this case, combined addition of NP would have offset the negative impact of single N addition. Thus, the addition of P does not seem to be a suitable strategy to enhance symbiotic N_2 fixation in a scenario of anthropogenic N enrichment of grasslands.

We observed no significant difference in the symbiotic N_2 fixation per area between the sites with annual or perennial legumes, because the higher biomass at sites dominated by annual legumes was counterbalanced by the higher %Ndfa of the sites dominated by perennials, irrespective of treatment (Fig. 4, Figure S3). This shows that there are no substantial differences in N_2 fixation on an area basis between grasslands dominated by annual and perennial legumes.

Across treatments, we found relatively low symbiotic N_2 fixation per area grassland ($3.4 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ in the control) compared to other studies (Carlsson and Huss-Danell 2003; Nyfeler et al. 2011; Peoples et al. 2012; Oberson et al. 2013). The reason for this seems to be our focus on natural and semi-natural grasslands with natural abundance of legumes (i.e. legumes were not deliberately introduced for the study), and the inclusion of some sites with low overall biomass production. The rate of N addition used in the present study ($100 \text{ kg N ha}^{-1} \text{ yr}^{-1}$) exceeds any present and even projected atmospheric N deposition levels (Ackerman et al. 2019). However, considering the common fertilization rates used in managed grasslands ranging from 20–30 up to $400 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ (Oenema et al. 2012; Klaus et al. 2018), the experimental rate used in our study resembles a realistic situation for many grasslands. The reduction of symbiotic N_2 fixation in grasslands by N addition increases the dependence of grassland biomass productivity on fertilization, which has several economic and environmental drawbacks (Lüscher et al. 2014).

Symbiotic N₂ fixation per unit legume biomass not affected by element addition

Although relatively high rates of N and P were added, we found no significant response of N₂ fixation per unit legume biomass to N, P or NP addition (Fig. 3B). This is due to the lack of element addition effect on legume N concentration and %Ndfa. Our finding is in disagreement with previous field studies in grasslands reporting a positive effect of P addition and a negative effect of N addition on %Ndfa (Høgh-Jensen et al. 2002; West et al. 2005; Burchill et al. 2014; Tzanakakis et al. 2017). The lack of response of N₂ fixation per unit legume biomass to P addition could indicate that symbiotic N₂ fixation was not limited by P in the majority of grasslands included in the present study, as P addition should increase N₂ fixation per unit biomass under strong P limitation due to the high ATP requirements of N₂ fixation (Almeida et al. 2000; Høgh-Jensen et al. 2002; Edwards et al. 2006). Similarly, the lack of differences between N and NP addition on N₂ fixation per unit biomass indicates that N addition did not induce a P limitation of N₂ fixation as previously described in pot experiments or tree plantations (Leidi and Rodríguez-Navarro 2000; Zheng et al. 2016). Otherwise, combined NP would have increased the N₂ fixation per unit biomass compared to single N addition. The reason why N addition did not cause a P limitation of N₂ fixation is likely that soil P availability is relatively high since the sites are located in the temperate zone which is dominated by relatively young soils (Figure S1).

The lack of response of N₂ fixation per unit biomass to N addition contrasts with previous results (Carlsson and Huss-Danell 2003) including experiments using urea as N source (Burchill et al. 2014), as in our study. We speculate that N addition had no significant effect on N₂ fixation per unit biomass because grasses and forbs were N limited (as indicated by the low N:P ratio of grasses and forbs in the control treatment), and their efficient uptake of additional N reduced the availability of added N to legumes, as described in previous studies (Nyfeler et al. 2011; Peoples et al. 2012; Oberson et al. 2013). In contrast, in other studies where legumes were deliberately introduced leading to a higher legume abundance (West et al. 2005; Burchill et al. 2014), non-fixing plants may not take up the added N which, in turn, could reduce the N₂ fixation per unit biomass. Another explanation might be that most legume species are permanent, rather than facultative

N₂ fixers and cannot shift their N source in spite of increased soil N availability (Menge and Hedin 2009).

We observed a higher N₂ fixation per unit biomass in the grasslands dominated by perennial legumes compared to the sites dominated by annual legumes. The reason for this could be that perennials can build up a symbiosis with N₂ fixation that last for several years, whereas annuals have to establish a new symbiosis with N₂ fixing microorganisms every year which makes this symbiosis likely less effective (Primieri et al. 2022). The differences in N₂ fixation per unit biomass between annual and perennial legume sites disappeared in the N treatment (Fig. 4D). This finding suggests that N addition has a very similar effect on N₂ fixation per unit biomass irrespective of whether the grassland is dominated by annual or perennial legumes.

Conclusions

Our results show that soil N enrichment by anthropogenic activities significantly reduces N₂ fixation in grasslands, and these effects cannot be reversed by additional P amendment. The negative effect of N addition on symbiotic N₂ fixation by legumes per area was caused exclusively by the negative effect of N addition on legume biomass, and not by an effect on the N₂ fixation rate per unit biomass, indicating that the legumes did not alter the rate at which they fixed N₂ per unit biomass. This reduction in symbiotic N₂ fixation per area increases the dependence of grassland productivity on fertilization and can ultimately change the ecological functioning of grasslands, affecting their net primary productivity as well as their above and belowground biodiversity, forage quality and provision of additional ecosystem services. Further, the unique global design of this study allowed us to derive an equation to correct for the effect of elevation on the isotope signature of N in grasses and non-fixing forbs which will be useful in future studies.

Acknowledgements We thank Renate Krauss and Fabrizio Protti for laboratory assistance and the Analytical Chemistry laboratory (CAN) of the Bayreuth Center of Ecological and Environmental Research (BayCEER) for performing parts of the chemical analyses. In addition, we thank the stable isotope laboratory of BayCEER for performing the isotope analyses. E.V. would like to thank Nikola Teutschová for her comments on a previous version of this manuscript. MCC and MNB thank to Companhia das Lezírias for granting access to the study site.

Authors' contributions EV, PMS and MSp conceived the ideas and designed methodology; EV, PMS and MSp collected the data; EV, PMS and MSp analyzed the data; EV, PMS and MSp led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

Funding Open access funding provided by Swedish University of Agricultural Sciences. EV, PMS and MS have been funded by the German Research Foundation through the Emmy Noether-program (grant SP1389/6–1). This work was generated using data from the Nutrient Network (NutNet, <https://nutnet.org>) experiment, funded at the site-scale by individual researchers. Coordination and data management have been supported by funding to E. Borer and E. Seabloom from the National Science Foundation Research Coordination Network (NSF-DEB-1042132) and Long-Term Ecological Research (NSF-DEB-1234162 and NSF-DEB-1831944 to Cedar Creek LTER) programs, and the Institute on the Environment (DG-0001–13). We also thank the Minnesota Supercomputer Institute for hosting project data and the Institute on the Environment for hosting Network meetings. AJ acknowledge the German Ministry for Education and Research (BMBF) for funding this research within the Biodiversa project SUSALPS; grant number: FKZ 031B0516C. NE and SH acknowledge support from the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, funded by the German Research Foundation (FZT 118). MNB also acknowledges funding through FCT contract DL 57/2016/CP1382/CT0030 and the COMPETE program through FEDER and FCT funding UID/BIA/50027/2013 and POCI-01–0145-FEDER-006821 and CEF by UID/AGR/00239/2019.

Declarations

Competing interest The authors have no relevant financial or non-financial interests to disclose.

Data availability Data will be publicly available on the Environmental Data Initiative platform.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Ackerman D, Millet DB, Chen X (2019) Global Estimates of Inorganic Nitrogen Deposition Across Four Decades. *Global Biogeochem Cycles* 33:100–107. <https://doi.org/10.1029/2018GB005990>
- Almeida JPF, Hartwig UA, Frehner M et al (2000) Evidence that P deficiency induces N feedback regulation of symbiotic N₂ fixation in white clover (*Trifolium repens* L.). *J Exp Bot* 51:1289–1297. <https://doi.org/10.1093/jxb/51.348.1289>
- Amarger N, Mariotti A, Mariotti F et al (1979) Estimate of symbiotically fixed nitrogen in field grown soybeans using variations in ¹⁵N Natural abundance. *Plant Soil* 52:269–280. <https://doi.org/10.1007/BF02184565>
- Amundson R, Austin AT, Schuur EAG et al (2003) Global patterns of the isotopic composition of soil and plant nitrogen. *Global Biogeochem Cycles* 17. <https://doi.org/10.1029/2002GB001903>
- Borer ET, Harpole WS, Adler PB et al (2014) Finding generality in ecology: A model for globally distributed experiments. *Methods Ecol Evol* 5:65–73. <https://doi.org/10.1111/2041-210X.12125>
- Borer ET, Grace JB, Harpole WS et al (2017) A decade of insights into grassland ecosystem responses to global environmental change. *Nat Ecol Evol* 1:1–7. <https://doi.org/10.1038/s41559-017-0118>
- Burchill W, James EK, Li D et al (2014) Comparisons of biological nitrogen fixation in association with white clover (*Trifolium repens* L.) under four fertiliser nitrogen inputs as measured using two ¹⁵N techniques. *Plant Soil* 385:287–302. <https://doi.org/10.1007/s11104-014-2199-1>
- Carlsson G, Huss-Danell K (2003) Nitrogen fixation in perennial forage legumes in the field. *Plant Soil* 253:353–372. <https://doi.org/10.1023/A:1024847017371>
- Choi W, Kwak J, Lim S et al (2017) Synthetic fertilizer and livestock manure differently affect $\delta^{15}\text{N}$ in the agricultural landscape: A review. *Agr Ecosyst Environ* 237:1–15. <https://doi.org/10.1016/j.agee.2016.12.020>
- Craine JM, Elmore AJ, Aida MPM et al (2009) Global patterns of foliar nitrogen isotopes and their relationships with climate, mycorrhizal fungi, foliar nutrient concentrations, and nitrogen availability. *New Phytol*
- Craine JM, Lee WG (2003) Covariation in leaf and root traits for native and non-native grasses along an altitudinal gradient in New Zealand. *Oecologia* 134:471–478. <https://doi.org/10.1007/s00442-002-1155-6>
- Edwards EJ, McCaffery S, Evans JR (2006) Phosphorus availability and elevated CO₂ affect biological nitrogen fixation and nutrient fluxes in a clover-dominated sward. *New Phytol* 169:157–167. <https://doi.org/10.1111/j.1469-8137.2005.01568.x>
- Gehring C, Vlek PLG (2004) Limitations of the ¹⁵N natural abundance method for estimating biological nitrogen fixation in Amazonian forest legumes. *Basic Appl Ecol* 5:567–580. <https://doi.org/10.1016/j.baae.2004.09.005>
- Güsewell S (2004) N:P ratios in terrestrial plants: Variation and functional significance. *New Phytol* 164:243–266. <https://doi.org/10.1111/j.1469-8137.2004.01192.x>

- Hansen JP, Vinther FP (2001) Spatial variability of symbiotic N₂ fixation in grass-white clover pastures estimated by the 15N isotope dilution method and the natural 15N abundance method. *Plant Soil* 230:257–266. <https://doi.org/10.1023/A:1010390901845>
- Hijmans RJ, Cameron SE, Parra JL et al (2005) Very high resolution interpolated climate surfaces for global land areas. *Int J Climatol* 25:1965–1978. <https://doi.org/10.1002/joc.1276>
- Hoegberg P (1997) Tansley Review No. 95 15N natural abundance in soil-plant systems. *New Phytol* 137(179):203. <https://doi.org/10.1046/j.1469-8137.1997.00808.x>
- Høgh-Jensen H, Schjoerring JK, Soussana JF (2002) The influence of phosphorus deficiency on growth and nitrogen fixation of white clover plants. *Ann Bot* 90:745–753. <https://doi.org/10.1093/aob/mcf260>
- Houlton BZ, Wang YP, Vitousek PM, Field CB (2008) A unifying framework for dinitrogen fixation in the terrestrial biosphere. *Nature* 454:327–330. <https://doi.org/10.1038/nature07028>
- Huber E, Wanek W, Gottfried M et al (2007) Shift in soil-plant nitrogen dynamics of an alpine-nival ecotone. *Plant Soil* 301:65–76. <https://doi.org/10.1007/s11104-007-9422-2>
- Hungria M, Vargas MAT (2000) Environmental factors affecting N₂ fixation in grain legumes in the tropics with an emphasis on Brazil. *Field Crops Res* 65:151–164. [https://doi.org/10.1016/S0378-4290\(99\)00084-2](https://doi.org/10.1016/S0378-4290(99)00084-2)
- Jacot KA, Lüscher A, Nösberger J, Hartwig UA (2000) Symbiotic N₂ fixation of various legume species along an altitudinal gradient in the Swiss Alps. *Soil Biol Biochem* 32:1043–1052. [https://doi.org/10.1016/S0038-0717\(00\)00012-2](https://doi.org/10.1016/S0038-0717(00)00012-2)
- Klaus VH, Kleinebecker T, Busch V et al (2018) Land use intensity, rather than plant species richness, affects the leaching risk of multiple nutrients from permanent grasslands. *Glob Change Biol* 24:2828–2840. <https://doi.org/10.1111/gcb.14123>
- Lamarque P, Tappeiner U, Turner C et al (2011) Stakeholder perceptions of grassland ecosystem services in relation to knowledge on soil fertility and biodiversity. *Reg Environ Change* 11:791–804. <https://doi.org/10.1007/s10113-011-0214-0>
- Leidi EO, Rodríguez-Navarro DN (2000) Nitrogen and phosphorus availability limit N₂ fixation in bean. *New Phytol* 147:337–346. <https://doi.org/10.1046/j.1469-8137.2000.00703.x>
- Lüscher A, Mueller-Harvey I, Soussana JF et al (2014) Potential of legume-based grassland-livestock systems in Europe: A review. *Grass Forage Sci* 69:206–228. <https://doi.org/10.1111/gfs.12124>
- McKey D (1994) Legumes and nitrogen: The evolutionary ecology of a nitrogen-demanding lifestyle. In: Sprent JI, McKey D (eds) *Advances in Legume Systematics 5: the Nitrogen Factor*. Royal Botanic Gardens, Kew, pp 211–228
- Menge DNL, Hedin LO (2009) Nitrogen fixation in different biogeochemical niches along a 120 000-year chronosequence in New Zealand
- Nuruzzaman M, Lambers H, Bolland MDA, Veneklaas EJ (2006) Distribution of carboxylates and acid phosphatase and depletion of different phosphorus fractions in the rhizosphere of a cereal and three grain legumes. *Plant Soil* 281:109–120. <https://doi.org/10.1007/s11104-005-3936-2>
- Nyfelner D, Huguenin-Elie O, Suter M et al (2011) Grass-legume mixtures can yield more nitrogen than legume pure stands due to mutual stimulation of nitrogen uptake from symbiotic and non-symbiotic sources. *Agr Ecosyst Environ* 140:155–163. <https://doi.org/10.1016/j.agee.2010.11.022>
- Oberson A, Frossard E, Bühlmann C et al (2013) Nitrogen fixation and transfer in grass-clover leys under organic and conventional cropping systems. *Plant Soil* 371:237–255. <https://doi.org/10.1007/s11104-013-1666-4>
- Oenema J, van Ittersum M, van Keulen H (2012) Improving nitrogen management on grassland on commercial pilot dairy farms in the Netherlands. *Agr Ecosyst Environ* 162:116–126. <https://doi.org/10.1016/j.agee.2012.08.012>
- Peñuelas J, Poulter B, Sardans J et al (2013) Human-induced nitrogen-phosphorus imbalances alter natural and managed ecosystems across the globe. *Nat Commun* 4:1–10. <https://doi.org/10.1038/ncomms3934>
- Peoples MB, Brockwell J, Hunt JR et al (2012) Factors affecting the potential contributions of N₂ fixation by legumes in Australian pasture systems. *Crop Pasture Sci* 63:759–786. <https://doi.org/10.1071/CP12123>
- Primieri S, Magnoli SM, Koffel T et al (2022) Perennial, but not annual legumes synergistically benefit from infection with arbuscular mycorrhizal fungi and rhizobia: a meta-analysis. *New Phytol* 233:505–514. <https://doi.org/10.1111/NPH.17787>
- Roscher C, Thein S, Weigelt A et al (2011) N₂ fixation and performance of 12 legume species in a 6-year grassland biodiversity experiment. *Plant Soil* 341:333–348. <https://doi.org/10.1007/s11104-010-0647-0>
- Schleuss PM, Widdig M, Biederman LA et al (2021) Microbial substrate stoichiometry governs nutrient effects on nitrogen cycling in grassland soils. *Soil Biol Biochem* 155. <https://doi.org/10.1016/j.soilbio.2021.108168>
- Soussana JF, Tallec T (2010) Can we understand and predict the regulation of biological N₂ fixation in grassland ecosystems? *Nutr Cycl Agroecosyst* 88:197–213. <https://doi.org/10.1007/s10705-009-9335-y>
- Stevens CJ, Dise NB, Mountford JO, Gowing DJ (2004) Impact of Nitrogen Deposition on the Species Richness of Grasslands. *Science* 303:1876–1879. <https://doi.org/10.1126/science.1094678>
- Suding KN, Collins SL, Gough L et al (2005) Functional- and abundance-based mechanisms explain diversity loss due to N fertilization. *Proc Natl Acad Sci U S A* 102:4387–4392. <https://doi.org/10.1073/pnas.0408648102>
- Suter M, Connolly J, Finn JA et al (2015) Nitrogen yield advantage from grass-legume mixtures is robust over a wide range of legume proportions and environmental conditions. *Glob Change Biol* 21:2424–2438. <https://doi.org/10.1111/gcb.12880>
- Tognetti PM, Prober SM, Báez S et al (2021) Negative effects of nitrogen override positive effects of phosphorus on grassland legumes worldwide. *Proc Natl Acad Sci U S A* 118. <https://doi.org/10.1073/pnas.2023718118>
- Tzanakakis V, Sturite I, Dörsch P (2017) Biological nitrogen fixation and transfer in a high latitude grass-clover

- grassland under different management practices. *Plant Soil* 421:107–122. <https://doi.org/10.1007/s11104-017-3435-2>
- Valentine AJ, Kleinert A, Benedito VA (2017) Adaptive strategies for nitrogen metabolism in phosphate deficient legume nodules. *Plant Sci* 256:46–52. <https://doi.org/10.1016/j.plantsci.2016.12.010>
- Vitousek PM, Shearer G, Kohl DH (1989) Foliar 15N natural abundance in Hawaiian rainforest: patterns and possible mechanisms. *Oecologia* 78:383–388. <https://doi.org/10.1007/BF00379113>
- West JB, HilleRisLambers J, Lee TD et al (2005) Legume species identity and soil nitrogen supply determine symbiotic nitrogen-fixation responses to elevated atmospheric [CO₂]. *New Phytol* 167:523–530. <https://doi.org/10.1111/j.1469-8137.2005.01444.x>
- Zheng M, Li D, Lu X et al (2016) Effects of phosphorus addition with and without nitrogen addition on biological nitrogen fixation in tropical legume and non-legume tree plantations. *Biogeochemistry* 131:65–76. <https://doi.org/10.1007/s10533-016-0265-x>
- Zheng M, Zhou Z, Luo Y et al (2019) Global pattern and controls of biological nitrogen fixation under nutrient enrichment: A meta-analysis. *Glob Change Biol* 25:3018–3030. <https://doi.org/10.1111/gcb.14705>
- Zhou Y, Cheng X, Fan J, Harris W (2016) Patterns and controls of foliar nitrogen isotope composition on the Qinghai-Tibet Plateau, China. *Plant Soil* 406:265–276. <https://doi.org/10.1007/s11104-016-2882-5>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.